



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Solution by the PROPOSER.

Let the equation of the parabola be $y^2=4px$, and that of the ellipse

$$\frac{(y-\beta)^2}{b^2} + \frac{(x-a)^2}{a^2} = 1.$$

A normal to the ellipse expressed by the tangent of the angle it makes with the axis of x has the equation

$$y-\beta = m(x-a) - \frac{(a^2-b^2)m}{\sqrt{(b^2m^2+a^2)}},$$

and in the case of the parabola $y=mx-2pm-pm^3$. Since the shortest distance is measured off on a common normal, the two equations should be identical, and therefore

$$\beta - ma - \frac{(a^2-b^2)m}{\sqrt{(b^2m^2+a^2)}} = -2pm - pm^3.$$

From this equation m is to be found. It leads to an equation of the 8th degree, which for numerical values presents no difficulty.

Substituting now the equation of the normal in both the equations of the parabola and ellipse, we find the co-ordinates of the points of intersection. Denoting these by x', y' , and x'', y'' , we find for the shortest distance the expression $\sqrt{[(x'-x'')^2 + (y'-y'')^2]}$.

The numerical equations given having a common value, x lying between 6 and 7, and intersecting, therefore present no suitable example.

Also solved by G. B. M. ZERR.

CALCULUS.

129. Proposed by JOHN M. COLAW, A. M., Monterey, Va.

Among all quadrilaterals inscribed in an ellipse, to determine that which contains the greatest area.

Solution by J. SCHEFFER, A. M., Hagerstown, Md.

Walton furnishes a solution of this interesting problem, which for its elegance and simplicity I reproduce here.

Let the equation of the ellipse be $x^2/a^2 + y^2/b^2 = 1$, and let the angular points of the quadrilateral be (x_1, y_1) , (x_2, y_2) , (x_3, y_3) , (x_4, y_4) . Then, u denoting the area of the quadrilateral,

$$2u = x_2y_1 - x_1y_2 + x_3y_2 - x_2y_3 + x_4y_3 - x_3y_4 + x_1y_4 - x_4y_1.$$

$$\frac{x_1^2}{a^2} + \frac{y_1^2}{b^2} = 1, \quad \frac{x_2^2}{a^2} + \frac{y_2^2}{b^2} = 1, \quad \frac{x_3^2}{a^2} + \frac{y_3^2}{b^2} = 1, \quad \frac{x_4^2}{a^2} + \frac{y_4^2}{b^2} = 1.$$

In order that u may be a maximum, we have, differentiating the last four equations, putting $du=0$, and using the method of indeterminate multipliers,

$$\frac{\lambda_1 x_1}{a^2} + y_4 - y_2 = 0, \quad \frac{\lambda_1 y_1}{b^2} - x_4 + x_2 = 0.$$

$$\text{Hence we have } \frac{x_1}{a^2}(x_4 - x_2) = -\frac{y_1}{b^2}(y_4 - y_2) \dots (1),$$

$$\frac{x_2}{a^2}(x_1 - x_3) = -\frac{y_2}{b^2}(y_1 - y_3) \dots (2), \quad \frac{x_3}{a^2}(x_2 - x_4) = -\frac{y_3}{b^2}(y_1 - y_4) \dots (3),$$

$$\frac{x_4}{a^2}(x_3 - x_1) = -\frac{y_4}{b^2}(y_3 - y_1) \dots (4).$$

$$\text{From (1) and (3) we have } \frac{y_3}{x_3} = \frac{y_1}{x_1} \dots (5), \text{ and from (2) and (4),}$$

$$\frac{y_4}{x_4} = \frac{y_2}{x_2} \dots (6). \quad \text{Also from (1) and (2), } \frac{1}{a^2}(x_1 x_4 - x_2 x_3) = -\frac{1}{b^2}(y_1 y_4 - y_2 y_3).$$

From the last three equations we see that

$$\frac{1}{a^2}(x_1 x_4 - x_2 x_3) = -\frac{1}{b^2} y_1 y_2 \left[\frac{x_4}{x_2} - \frac{x_3}{x_1} \right], \text{ and therefore } \frac{y_1 y_2}{x_1 x_2} = -\frac{b^2}{a^2} \dots (7).$$

The equations (5) and (6) show that the diagonals of the quadrilaterals are diameters of the ellipse, and (7) shows that they are conjugate diameters.

Also solved by *G. B. M. ZERR*, *A. H. HOLMES*, and *L. C. WALKER*.

130. Proposed by *J. SCHEFFER*, *A. M.*, Hagerstown, Md.

$$\text{Solve the differential equation } x^x \left(\frac{dy}{dx} + y \log x \right) - a = 0.$$

I. Solution by *W. E. HEAL*, Marion, Ind.

Writing the equation $\frac{dy}{dx} + y \log x = a/x^x$, we have a linear equation of the standard form, and by the well known formula, $y = -Ca/x^x$.

II. Solution by *C. HORNUNG*, *A. M.*, Professor of Mathematics, Heidelberg University, Tiffin, Ohio.

Dividing by x^x and transposing we get $dy/dx + \log x \cdot y = ax^{-x}$, the regular form of the linear equation.

Using Bernoulli's method, we proceed as follows:

$$\text{Let } y = uz; \text{ then } dy/dx = u \frac{dz}{dx} + z \frac{du}{dx}, \text{ and the given equation becomes}$$

$$u \frac{dz}{dx} + z \left(\frac{du}{dx} + u \log x \right) = ax^{-x}.$$